



[cancer.org](https://www.cancer.org) | 1.800.227.2345

Immunotherapy

Immunotherapy is treatment that uses a person's own immune system to fight cancer. Immunotherapy can boost or change how the immune system works so it can find and attack cancer cells. If your treatment plan includes immunotherapy, knowing how it works and what to expect can often help you prepare for treatment and make informed decisions about your care.

[How Immunotherapy Is Used to Treat Cancer](#)

[Monoclonal Antibodies](#)

[CAR T-cell Therapy](#)

[Immune Checkpoint Inhibitors](#)

[Cancer Vaccines](#)

[Cytokines](#)

[Immunomodulators](#)

[Immunotherapy Safety](#)

Related Topics

- [Intravenous \(IV\) Lines, Catheters, and Ports Used in Cancer Treatment](#)
- [Tubes, Catheters, and Drains Used in Cancer Treatment and Care](#)
- [Compassionate Drug Use](#)
- [Off-label Drug Use](#)

Video

This short video provides answers to some of the most common questions that people with cancer have about immunotherapy.

[Watch this video on YouTube](#)

as “foreign” and attacks them. The immune response can destroy anything containing the foreign substance, such as germs or cancer cells.

The immune system has a tougher time targeting cancer cells, though. This is because cancer starts when normal, healthy cells become changed or altered and start to grow out of control. Because cancer cells actually start in normal cells, the immune system doesn't always recognize them as foreign.

Clearly there are limits on the immune system's ability to fight cancer on its own, because many people with healthy immune systems still develop cancer:

- Sometimes the immune system doesn't see the cancer cells as foreign because the cells aren't different enough from normal cells.
- Sometimes the immune system recognizes the cancer cells, but the response might not be strong enough to destroy the cancer.
- Cancer cells themselves can also give off substances that keep the immune system from finding and attacking them.

To overcome this, researchers have found ways to help the immune system recognize cancer cells and strengthen its response so that it will destroy them. In this way, your own body is actually getting rid of the cancer, with some help from science.

Types of cancer immunotherapy

There are several main types of immunotherapy used to treat cancer, and many are being studied. **For more information about immunotherapy as a treatment for a specific cancer, please [choose a cancer type](#)¹.**

- **Checkpoint inhibitors:** These drugs basically take the ‘brakes’ off the immune system, which helps it recognize and attack cancer cells.
- **Chimeric antigen receptor (CAR) T-cell therapy:** This therapy takes some T-cells from a patient's blood, mixes them with a special virus that makes the T-cells learn how to attach to tumor cells, and then gives the cells back to the patient so they can find, attach to, and kill the cancer.
- **Cytokines:** This treatment uses *cytokines* (small proteins that carry messages between cells) to stimulate the immune cells to attack cancer.
- **Immunomodulators:** This group of drugs generally boosts parts of the immune system to treat certain types of cancer.
- **Cancer vaccines:** Vaccines are substances put into the body to start an immune

response against certain diseases. We usually think of them as being given to healthy people to help prevent infections. But some vaccines can help prevent or treat cancer.

- **Monoclonal antibodies (mAbs or MoAbs):** These are man-made versions of immune system proteins. mAbs can be very useful in treating cancer because they can be designed to attack a very specific part of a cancer cell.
- **Oncolytic viruses:** This treatment uses viruses that have been modified in a lab to infect and kill certain tumor cells.

DeSelm CJ, Tano ZE, Varghese AM, et al. CAR T-cell therapy for pancreatic cancer. *J Surg Oncol*. 2017; 16(1):63-74.

Gatti-Mays ME, Redman JM, Collins JM, et al. Cancer vaccines: Enhanced immunogenic modulation through therapeutic combinations. *Hum Vaccin Immunother*. 2017; 13(11):2561-2574. Accessed at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5703410/> on December 19, 2019.

Hafeez U, Gan HK, Scott AM. Monoclonal antibodies as immunomodulatory therapy against cancer and autoimmune diseases. *Curr Opin Pharmacol*. 2018; 41:114-121.

Hill JA, Giralt S, Torgerson TR, et al. CAR-T- and a side order of IgG, to go?- Immunoglobulin replacement in patients receiving CAR-T cell therapy. *Blood Rev*. 2019 [Accepted manuscript]. Accessed at <https://www.ncbi.nlm.nih.gov/pubmed/31416717> on December 19, 2019.

Ling DC, Bakkenist CJ, Ferris RL et al. Role of immunotherapy in head and neck cancer. *Semin Radiat Oncol*. 2018; 28(1): 12-16.

Maeng H, Terabe M, Berzofsky JA. Cancer vaccines: Translation from mice to human clinical trials. *Curr Opin Immunol*. 2018; 51:111-122. Accessed at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5943163/> on December 19, 2019.

Mestermann K, Giavridis T, Weber J, , et al. The tyrosine kinase inhibitor dasatinib acts as a pharmacologic on/off switch for CAR T cells. *Sci Transl Med*. [Abstract]. 2019; 11(499).

Myers DR, Wheeler B, Roose JP. mTOR and other effector kinase signals that impact T cell function and activity. *Immunol Rev*. 2019; 291(1):134-153.

National Cancer Institute (NCI). *CAR T cells: Engineering patients' immune cells to treat their cancers*. Accessed at <https://www.cancer.gov/about-cancer/treatment/research/car-t-cells> on December 19, 2019.

National Cancer Institute (NCI). *Immunotherapy to treat cancer*. Accessed at <https://www.cancer.gov/about-cancer/treatment/types/immunotherapy> on December 19, 2019.

Russell SJ, Barber GN. Oncolytic viruses as antigen-agnostic cancer vaccines. *Cancer Cell*. 2018;33(4): 599-605. Accessed at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5918693/> on December 19, 2019.

Wraith DC. The future of immunotherapy: A 20-year perspective. *Front Immunol*. 2017;8:1668. Accessed at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5712390/> on December 19, 2019.

Last Revised: December 27, 2019

Monoclonal Antibodies and Their Side Effects

One way the immune system protects the body from germs and other foreign substances is by making large numbers of **antibodies**. An antibody is a protein that sticks to a specific protein called an **antigen**. Antibodies circulate throughout the body until they find and attach to the antigen. Once attached, they can help other parts of the immune system to destroy the cells containing the antigen.

Researchers can design antibodies that specifically target a certain antigen, such as one found on cancer cells. They can then make many copies of that antibody in the lab. These are known as *monoclonal antibodies* (mAbs or Moabs).

- [What mAbs are made of](#)
- [Types of mAbs used to treat cancer](#)
- [Possible side effects of monoclonal antibodies](#)

Monoclonal antibodies are used to treat many diseases, including some types of cancer. To make a monoclonal antibody, researchers first have to identify the right antigen to attack. Finding the right antigens for cancer cells is not always easy, and so far mAbs have proven to be more useful against some cancers than others.

NOTE: Some monoclonal antibodies used to treat cancer are sometimes considered to be a type of [targeted therapy](#)¹ because they work by attaching to a specific target on a cancer cell and stopping it from functioning. But other monoclonal antibodies act like immunotherapy because they help the immune system find and attack cancer cells more effectively.

What mAbs are made of

Monoclonal antibodies are man-made proteins that act like human antibodies in the immune system. There are 4 different ways they can be made and are named based on what they are made of.

- **Murine:** These are made from mouse proteins, and the names of the treatments end in -omab.
- **Chimeric:** These proteins are a combination of part mouse and part human, and the names of the treatments end in -ximab.
- **Humanized:** These are made from small parts of mouse proteins attached to human proteins, and the names of the treatments end in -zumab
- **Human:** These are fully human proteins, and the names of the treatments end in -umab.

Types of mAbs used to treat cancer

Naked monoclonal antibodies

Naked mAbs are antibodies that have no drug or radioactive material attached to them. They work by themselves. These are the most common type of mAbs used to treat cancer. Most naked mAbs attach to antigens on cancer cells, but some work by binding to antigens on other, non-cancerous cells, or even free-floating proteins. Naked mAbs can work in different ways.

- Some boost a person's immune response against cancer cells by attaching to them and acting as a marker for the body's immune system to destroy them. An example is rituximab (Rituxan), which is used to treat some types of non-Hodgkin lymphoma (NHL) and other cancers. Rituximab binds to the CD antigen, which is found on cells called *B lymphocytes* (which include some types of lymphoma cells). Once attached, the antibody attracts immune cells to destroy these cells.
- Some naked mAbs boost the immune response by targeting immune system checkpoints. (See [Immune Checkpoint Inhibitors and Their Side Effects](#).)
- Other naked mAbs work mainly by attaching to and blocking proteins on cancer cells (or other nearby cells) that help cancer cells grow or spread. For example, trastuzumab (Herceptin) is an antibody against the HER2 protein. Breast and stomach cancer cells sometimes have large amounts of this protein on their surface, which helps these cells grow. Trastuzumab binds to these proteins and stops them from becoming active.

Conjugated monoclonal antibodies

Conjugated mAbs are connected to a chemotherapy drug or a radioactive particle. These mAbs are used as a homing device to take one of these substances directly to the cancer cells. The mAb circulates throughout the body until it can find and hook onto the target antigen. It then delivers the toxic substance where it is needed most. Conjugated mAbs are also sometimes referred to as *tagged*, *labeled*, or *loaded* antibodies.e.

which is something like an allergic reaction. This is more common while the drug is first being given. Possible symptoms can include:

- Fever
- Chills
- Weakness
- Headache
- Nausea
- Vomiting
- Diarrhea
- Low blood pressure
- Rashes

Compared with chemotherapy drugs, naked mAbs tend to have fewer serious side effects. But they can still cause problems in some people.

Some mAbs can have side effects that are related to the antigens they target. For example:

- Bevacizumab (Avastin) is an mAb that targets a protein called *VEGF* that affects tumor blood vessel growth. It can cause side effects such as high blood pressure, bleeding, poor wound healing, blood clots, and kidney damage.
- Cetuximab (Erbix) is an antibody that targets a cell protein called *EGFR*, which is found on normal skin cells (as well as some types of cancer cells). This drug can cause serious rashes in some people.

Hyperlinks

1. www.cancer.org/cancer/managing-cancer/treatment-types/targeted-therapy.html

References

Acrotech Biopharma. *What is Zevalin?* Accessed at <http://www.zevalin.com/patient/is-zevalin-right-for-you/what-is-zevalin> on December 19, 2019.

CAR T-cell Therapy and Its Side Effects

- [Approved CAR T-cell therapies](#)
- [Possible CAR T-cell therapy side effects](#)

Chimeric antigen receptor (CAR) T-cell therapy is a way to get immune cells called *T cells* (a type of white blood cell) to fight cancer by changing them in the lab so they can find and destroy cancer cells. CAR T-cell therapy is also sometimes talked about as a type of *cell-based gene therapy*, because it involves altering the genes inside T cells to help them attack the cancer.

This type of treatment can be very helpful in treating some types of cancer, even when other treatments are no longer working.

How CAR T-cell therapy works

Immune receptors and foreign antigens

The immune system recognizes foreign substances in the body by finding proteins called *antigens* on the surface of those cells. Immune cells called *T cells* have their own proteins called *receptors* that attach to foreign antigens and help trigger other parts of the immune system to destroy the foreign substance.

The relationship between antigens and immune receptors is like a lock and key. Just as a lock can only be opened with the right key, each foreign antigen has a unique immune receptor that is able to bind to it.

Cancer cells also have antigens, but if your immune cells don't have the right receptors, they can't attach to the antigens and help destroy the cancer cells.

Chimeric antigen receptors (CARs)

In CAR T-cell therapies, T cells are taken from the patient's blood and are changed in the lab by adding a gene for a receptor (called a *chimeric antigen receptor* or *CAR*), which helps the T cells attach to a specific cancer cell antigen. The CAR T cells are then given back to the patient.

Since different cancers have different antigens, each CAR is made for a specific cancer's antigen. For example, in certain kinds of leukemia or lymphoma, the cancer cells have an antigen called CD19. The CAR T-cell therapies to treat these cancers are made to attach to the CD19 antigen and will not work for a cancer that does not have the CD19 antigen.

Getting CAR T-cell therapy

The process for CAR T-cell therapy can take several weeks.

Collecting the T cells

Examples of CAR T-cell therapies currently approved include:

- Tisagenlecleucel, also known as tisa-cel (Kymriah)
- Axicabtagene ciloleucel, also known as axi-cel (Yescarta)
- Brexucabtagene autoleucel, also known as brexu-cel (Tecartus)
- Lisocabtagene maraleucel, also known as liso-cel (Breyanzi)
- Idecabtagene vicleucel, also known as ide-cel (Abecma)
- Ciltacabtagene autoleucel, also known as cilta-cel (Carvykti)

Many other CAR T-cell therapies (and similar types of treatment) are now being studied in clinical trials, in the hope of treating other types of cancer as well.

Possible CAR T-cell therapy side effects

CAR T-cell therapy can be very effective against some types of hard-to-treat cancers, but it can also sometimes cause serious or even life-threatening side effects. Because of this, it needs to be given in a medical center that is specially trained in its use, and patients need to be watched closely for several weeks after getting the CAR T cells.

Cytokine release syndrome (CRS): As CAR T cells multiply, they can release large amounts of chemicals called *cytokines* into the blood, which can ramp up the immune system. Serious side effects from this release can include:

- High fever and chills
- Trouble breathing
- Severe nausea, vomiting, and/or diarrhea
- Feeling dizzy or lightheaded
- Headaches
- Fast heartbeat
- Feeling very tired
- Muscle and/or joint pain

As doctors are gaining more experience with CAR T-cell therapy, they are learning how to recognize CRS early, as well as how to treat it.

Nervous system problems: This treatment can sometimes have serious effects on the nervous system, which can result in symptoms such as:

- Headaches

- Changes in consciousness
- Confusion or agitation
- Seizures
- Shaking or twitching (tremors)
- Trouble speaking and understanding
- Loss of balance

Because of the risk of these side effects, adult patients are typically advised not to drive, operate heavy machinery, or do any other potentially dangerous activities for at least several weeks after getting treatment.

Other serious side effects: Other possible serious side effects of CAR T-cell therapy can include:

- Allergic reactions during the infusion
- Abnormal levels of minerals in the blood, such as low potassium, sodium, or phosphorous levels
- A weakened immune system, with an increased risk of serious infections
- Low blood cell counts, which can increase the risk of infections, fatigue, and bruising or bleeding

If you are getting CAR T-cell therapy, it's very important to report any side effects to your health care team right away, as there are often medicines that can help treat them.

Hyperlinksbruigs 1.5 Ts (l)Tj 0 Ts 0ew4 7 /F2 12 Tf 0 0 0 rg /GS633 rg /GS624 gs 1.

(ASCO). Understanding immunotherapy. Accessed at cancer.net. Content is no longer available.

Bayer VR, Davis ME, Gordan RA, et al. Immunotherapy. In Olsen MM, LeFebvre KB, Brassil KJ, eds. *Chemotherapy and Immunotherapy Guidelines and Recommendations for Practice*. Pittsburgh, PA: Oncology Nursing Society; 2019:149-189.

National Cancer Institute (NCI). (NCI). CAR T cells: Engineering patients' immune cells to treat their cancers. Accessed at <https://www.cancer.gov/about-cancer/treatment/research/car-t-cells> on December 19, 2019.

Porter DL, Maloney DG. Cytokine release syndrome (CRS). UpToDate. 2021. Accessed at <https://www.uptodate.com/contents/cytokine-release-syndrome-crs> on April 01, 2021.

Last Revised: March 1, 2022

Immune Checkpoint Inhibitors and Their Side Effects

An important part of the immune system is its ability to tell between normal cells in the body and those it sees as “foreign” (such as germs and cancer cells). This allows the immune system to attack the foreign cells while leaving normal cells alone.

Part of how the immune system does this is by using “checkpoint” proteins on immune cells. The checkpoints act like switches that need to be turned on (or off) to start an immune response. But cancer cells sometimes find ways to use these checkpoints to avoid being attacked by the immune system.

- [PD-1 and PD-L1 inhibitors](#)
- [CTLA-4 inhibitors](#)
- [LAG-3 inhibitors](#)
- [Side effects of checkpoint inhibitors](#)

Medicines known as [monoclonal antibodies](#) can be designed to target these checkpoint proteins. These drugs are called **immune checkpoint inhibitors**(or just **checkpoint inhibitors**).

Checkpoint inhibitors don't kill cancer cells directly. They work by helping the immune system to better find and attack the cancer cells, wherever they are in the body.

Medicines that target different checkpoint proteins are now used to treat some types of cancer. All of these drugs are given as an infusion into a vein (IV).

PD-1 and PD-L1 inhibitors

PD-1 is a checkpoint protein on immune cells called *T cells*. It normally acts as a type of “off switch” that helps keep the T cells from attacking other cells in the body. It does this when it attaches to PD-L1, a protein on some normal (and cancer) cells. When PD-1 binds to PD-L1, it basically tells the T cell to leave the other cell alone. Some cancer cells have large amounts of PD-L1, which helps them hide from an immune attack.

Monoclonal antibodies that target either PD-1 or PD-L1 can block this binding and boost the immune response against cancer cells.

PD-1 inhibitors

Examples of drugs that target PD-1 include:

- **Pembrolizumab (Keytruda)**
- **Nivolumab (Opdivo)**
- **Cemiplimab (Libtayo)**

PD-L1 inhibitors

Examples of drugs that target PD-L1 include:

- **Atezolizumab (Tecentriq)**
- **Avelumab (Bavencio)**
- **Durvalumab (Imfinzi)**

Both PD-1 and PD-L1 inhibitors have been shown to be helpful in treating many different types of cancer.

CTLA-4 inhibitors

CTLA-4 is another checkpoint protein on some T cells that acts as a type of “off switch” to help keep the immune system in check.

Ipilimumab (Yervoy) and **tremelimumab (Imjuno)** are monoclonal antibodies that attach to CTLA-4 and stop it from working. This can help boost the body’s immune response against cancer cells.

These drugs are typically used along with a PD-1 or PD-L1 inhibitor. These combinations can be used to treat several types of cancer.

LAG-3 inhibitors

LAG-3 is a checkpoint protein on some types of immune cells that normally acts as a type of “off switch” to help keep the immune system in check.

Relatlimab is a monoclonal antibody that attaches to LAG-3 and stops it from working. This can help boost the body’s immune response against cancer cells.

This drug is given along with the PD-1 inhibitor nivolumab (in a combination known as **Opdualag**). It can be used to treat melanoma of the skin, and it’s being studied for use in several other types of cancer.

Side effects of checkpoint inhibitors

Some of the more common side effects of checkpoint inhibitors include:

- Diarrhea
- Fatigue
- Cough
- Nausea
- Skin rash
- Poor appetite
- Constipation
- Muscle and joint pain

Other, more serious side effects occur less often:

Infusion reactions: Some people might have an infusion reaction while getting these drugs. This is like an allergic reaction, and can include fever, chills, flushing of the face,

rash, itchy skin, feeling dizzy, wheezing, and trouble breathing. It's important to tell your doctor or nurse right away if you have any of these symptoms while getting one of these drugs.

Autoimmune reactions: By targeting a a3r

Cancer Vaccines and Their Side Effects

as measles and chicken pox. These vaccines use weakened or killed germs like viruses or bacteria to start an immune response in the body. Getting the immune system ready to defend against these germs helps keep people from getting infections.

- [Vaccines to help prevent cancer](#)
- [Vaccines to treat cancer](#)
- [Other vaccines](#)

Most vaccines used to treat cancer work the same way, but they make the person's immune system attack cancer cells. The goal is to help treat cancer or to help keep it from coming back after other treatments. But there are also some vaccines that may actually help prevent certain cancers.

Vaccines to help prevent cancer

Some cancers are caused by viruses. Vaccines that help protect against infections with these viruses might also help prevent some of these cancers.

- Some strains of the [human papillomavirus \(HPV\)](#)¹ have been linked to cervical, anal, throat, vaginal, vulvar, and penile cancers. In fact, most cervical cancers are caused by infection with HPV. Vaccinating children and certain young adults against HPV helps protect against cervical cancer and the other 5 cancers HPV can cause. Read more in [Protect Against HPV](#)².
- People who have chronic (long-term) infections with the hepatitis B virus (HBV) are at higher risk for [liver cancer](#)³. Getting the vaccine to help prevent HBV infection may lower some people's risk of getting liver cancer.

These are traditional preventive vaccines that target the viruses that can cause certain cancers. They may help protect against some cancers, but they don't target cancer cells directly because cancer cells have not yet been formed or found.

These types of vaccines are only useful for cancers known to be caused by infections. But most cancers, including colorectal, lung, prostate, and breast cancers, are not thought to be caused by infections.

Vaccines to treat cancer

Cancer treatment vaccines are different from the vaccines that work against viruses. These vaccines try to get the immune system to mount an attack against cancer cells in

the body. Instead of preventing disease, they are meant to get the immune system to attack a disease that already exists.

Some cancer treatment vaccines are made up of cancer cells, parts of cells, or pure antigens (certain proteins on the cancer cells). Sometimes a patient's own immune cells are removed and exposed to these substances in the lab to create the vaccine. Once the vaccine is ready, it's injected into the body to increase the immune response against cancer cells.

Vaccines are often combined with other substances or cells called *adjuvants* that help boost the immune response even further.

Cancer vaccines cause the immune system to attack cells with one or more specific antigens. Because the immune system has special cells for memory, it's hoped that the vaccine might continue to work long after it's given.

- **Sipuleucel-T (Provenge):** This drug is used to treat advanced prostate cancer that is no longer being helped by hormone therapy. [Side effects](#)⁴ are usually mild and can include fever, chills, fatigue, back and joint pain, nausea, and headache. A few men may have more severe symptoms, including problems breathing and high blood pressure.
- **Talimogene laherparepvec (T-VEC):** This vaccine is approved to treat advanced melanoma skin cancer. It is made from a herpes virus that has been altered in the lab to produce a substance that the body normally produces, called a *cytokine*. This cytokine boosts the immune system and can cause flu-like symptoms for a short time.

Other vaccines

Other types of cancer vaccines have shown some promise in clinical trials, but they are not yet approved in the U.S. to treat cancer.

Hyperlinks

1. www.cancer.org/cancer/risk-prevention/hpv.html

2. www.cancer.org/cancer/risk-prevention/hpv/hpv-vaccine.html
3. www.cancer.org/cancer/types/liver-cancer.html
4. www.cancer.org/cancer/managing-cancer/side-effects.html

References

American Society of Clinical Oncology (ASCO). ASCO Annual Meeting 2019: Immunotherapy for lung cancer, gastrointestinal cancers and targeted therapy for breast cancer. Accessed at cancer.net. Content is no longer available.

American Society of Clinical Oncology (ASCO). Understanding immunotherapy. Accessed at cancer.net. Content is no longer available.

Bayer VR, Davis ME, Gordan RA, et al. Immunotherapy. In Olsen MM, LeFebvre KB, Brassil KJ, eds. *Chemotherapy and Immunotherapy Guidelines and Recommendations for Practice*. Pittsburgh, PA: Oncology Nursing Society; 2019:149-189.

Coventry BJ. Therapeutic vaccination immunomodulation: Forming the basis of all cancer immunotherapy. *Ther Adv Vaccines Immunother*. 2019; 1:7:2515135519862234. Accessed at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6676259/> on December 19, 2019.

DeMaria PJ, Bilusic M. Cancer vaccines. *Hematol Oncol Clin North Am*. 2019; 33(2):199-214.

Saslow D, Andrews KS, Manassaram-Baptiste D, et al. Human papillomavirus vaccination 2020 guideline update: American Cancer Society guideline adaptation. *CA Cancer J Clin*. 2020; DOI: 10.3322/caac.21616.

Last Revised: January 8, 2020

Cytokines and Their Side Effects

Cytokines are small proteins that are crucial in controlling the growth and activity of other immune system cells and blood cells. When released, they signal the immune system to do its job. Cytokines affect the growth of all blood cells and other cells that help the body's immune and inflammation responses. They also help to boost anti-cancer activity by sending signals that can help make abnormal cells die and normal cells live longer.

- [Interleukins](#)
- [Interferons](#)

One specific type of cytokine is called a *chemokine*. A chemokine can make immune cells move toward a target. There are different kinds of chemokines, including *interleukins*, *interferons*, *tumor necrosis factors*, and *growth factors*.

Some cytokines can be made in a lab and are used to treat cancer. Some are used to help prevent or manage chemotherapy side effects. They are injected, either under the skin, into a muscle, or into a vein. The most common ones are interleukins and interferons.

Interleukins

Interleukins are a group of cytokines that act as chemical signals between white blood cells. Interleukin-2 (IL-2) helps immune system cells grow and divide more quickly. A man-made version of IL-2 is approved to treat advanced [kidney cancer](#)¹ and metastatic [melanoma](#)². IL-2 can be used as a single drug treatment for these cancers, or it can be combined with chemotherapy or with other cytokines such as interferon-alfa.

[Side effects](#)³ of IL-2 can include flu-like symptoms such as chills, fever, fatigue, and confusion. Some have nausea, vomiting, or diarrhea. Many people develop low blood pressure, which can be treated with other medicines. Rare but potentially serious side effects include an abnormal heartbeat, chest pain, and other heart problems. Because of these possible side effects, if IL-2 is given in high doses, it must be done in a hospital.

Other interleukins, such as IL-7, IL-12, and IL-21, continue to be studied for use against cancer too, both as adjuvants and as stand-alone agents.

Interferons

Interferons are chemicals that help the body resist virus infections and cancers. The

types of interferon (IFN) are named after the first 3 letters of the Greek alphabet:

- IFN-alfa
- IFN-beta
- IFN-gamma

Only IFN-alfais used to treat cancer. It boosts the ability of certain immune cells to attack cancer cells. It may also slow the growth of cancer cells directly, as well as the blood vessels that tumors need to grow.

IFN-alfa can be used to treat these cancers:

- Hairy cell leukemia
- Chronic myelogenous leukemia (CML)
- Follicular non-Hodgkin lymphoma
- Cutaneous (skin) T-cell lymphoma
- Kidney cancer
- Melanoma
- Kaposi sarcoma

Side effects of interferons can include:

- Flu-like symptoms (chills, fever, headache, fatigue, loss of appetite, nausea, vomiting)
- Low white blood cell counts (which increase the risk of infection)
- Skin rashes
- Thinning hair

These side effects can be severe and can make treatment with interferon hard for many people to tolerate. Most side effects don't last long after the treatment stops, but fatigue can last longer. Other rare long-term effects include damage to nerves, including those in the brain and spinal cord.

Hyperlinks

1. www.cancer.org/cancer/types/kidney-cancer.html

2. www.cancer.org/cancer/types/melanoma-skin-cancer.html
3. www.cancer.org/cancer/managing-cancer/side-effects.html

References

American Society of Clinical Oncology (ASCO). ASCO Annual Meeting 2019: Immunotherapy for lung cancer, gastrointestinal cancers and targeted therapy for breast cancer. Accessed at cancer.net. Content is no longer available.

American Society of Clinical Oncology (ASCO). Understanding immunotherapy. Accessed at cancer.net. Content is no longer available.

Bayer VR, Davis ME, Gordan RA, et al. Immunotherapy. In Olsen MM, LeFebvre KB, Brassil KJ, eds. *Chemotherapy and Immunotherapy Guidelines and Recommendations for Practice*. Pittsburgh, PA: Oncology Nursing Society; 2019:149-189.

Brodsky AN. *Cancer immunotherapy: The year in review and a look at the year ahead*. Cancer Research Institute. Accessed at <https://www.cancerresearch.org/blog/january-2019/cancer-immunotherapy-2018-review-2019-predict> on December 19, 2019.

Hafeez U, Gan HK, Scott AM. Monoclonal antibodies as immunomodulatory therapy against cancer and autoimmune diseases. *Curr Opin Pharmacol*. 2018; 41:114-121.

Wraith DC. The future of immunotherapy: A 20-year perspective. *Front Immunol*. 2017;8:1668. Accessed at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5712390/> on December 19, 2019.

Last Revised: December 27, 2019

Immunomodulators and Their Side Effects

Immunomodulators are a group of drugs that mainly target the pathways that treat multiple myeloma and a few other cancers. They have many ways to work, including working on the immune system directly by turning down some proteins and turning up others.

- [Thalidomide, lenalidomide, and pomalidomide](#)
- [Bacillus Calmette-Guérin](#)
- [Imiquimod](#)

Thalidomide, lenalidomide, and pomalidomide

Thalidomide (Thalomid), lenalidomide (Revlimid), and pomalidomide (Pomalyst) are known as *immunomodulating drugs* (or IMiDs).

These drugs can cause side effects such as drowsiness, fatigue, constipation, low blood cell counts, and neuropathy (painful nerve damage). There is also an increased risk of serious blood clots (that start in the leg and can travel to the lungs). These tend to be more likely with thalidomide than with the other drugs.

These drugs can also cause severe birth defects if taken during pregnancy.

Bacillus Calmette-Guérin

Bacillus Calmette-Guérin (BCG) is a germ that doesn't cause serious disease in humans, but it does infect human tissues and helps activate the immune system. This makes BCG useful as a form of cancer immunotherapy. BCG was one of the earliest immunotherapies used against cancer and is still being used today.

BCG is used to treat early stage bladder cancer. It is a liquid put into the bladder through a catheter. BCG attracts the body's immune system cells to the bladder, where they can attack the bladder cancer cells. Treatment with BCG can cause symptoms that are like having the flu, such as fever, chills, and fatigue. It can also cause a burning feeling in the bladder. BCG can also be used to treat some melanoma skin cancers by injecting it directly into the tumors. It's also used as a vaccine against tuberculosis.

Imiquimod

Imiquimod is a drug that is applied to the skin as a cream. It stimulates a local immune response against skin cancer cells. It is used to treat some very early stage skin cancers (or pre-cancers), especially if they are in sensitive areas such as on the face.

The cream is applied anywhere from once a day to twice a week for several months. Some people have serious skin reactions to this drug.

References

American Society of Clinical Oncology (ASCO). ASCO Annual Meeting 2019: Immunotherapy for lung cancer, gastrointestinal cancers and targeted therapy for breast cancer. Accessed at cancer.net. Content is no longer available.

American Society of Clinical Oncology (ASCO). Understanding immunotherapy. Accessed at cancer.net. Content is no longer available.

Bayer VR, Davis ME, Gordan RA, et al. Immunotherapy. In Olsen MM, LeFebvre KB, Brassil KJ, eds. *Chemotherapy and Immunotherapy Guidelines and Recommendations for Practice*. Pittsburgh, PA: Oncology Nursing Society; 2019:149-189.

Brodsky AN. *Cancer immunotherapy: The year in review and a look at the year ahead*. Cancer Research Institute. Accessed at <https://www.cancerresearch.org/blog/january-2019/cancer-immunotherapy-2018-review-2019-predict> on December 19, 2019.

Hafeez U, Gan HK, Scott AM. Monoclonal antibodies as immunomodulatory therapy against cancer and autoimmune diseases. *Curr Opin Pharmacol*. 2018; 41:114-121.

Wraith DC. The future of immunotherapy: A 20-year perspective. *Front Immunol*. 2017;8:1668. Accessed at <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5712390/> on December 19, 2019.

Last Revised: December 27, 2019

Immunotherapy Safety

Much is known about the need to protect others from exposure to [traditional or standard](#) [cancer about to](#) [reg n pet.8 0 G 1 w 0 0](#). Osure to December 27, 2 Tf 0 rs (. Pittsburg g /F (Referwhoaha

taking the same precautions. This is especially true when immunotherapy drugs are given to treat cancer in combination with other drugs that are known to be hazardous, so your cancer care team will take precautions to protect themselves and others from exposure to them.

- [Precautions the cancer care team might take](#)
- [Special precautions when taking oral or topical immunotherapy](#)
- [Keeping family and friends safe](#)

Precautions the cancer care team might take

You may notice special clothing and protective equipment being worn by the nurses and other members of your cancer care team. Pharmacists and nurses who prepare drugs to treat cancer use a special type of pharmacy that must meet certain regulations. If you are being cared for in a treatment center, the nurses and others who give treatment and help take care of patients afterwards wear protective clothing, such as 2 pairs of special gloves and a gown, and sometimes goggles or a face shield. If you're getting immunotherapy through an IV, there might be a disposable pad under the infusion tubing to protect the surface of the bed or chair.

Special precautions when taking oral or topical immunotherapy

Oral immunotherapy that you take by mouth and swallow, or topical immunotherapy that you rub on your skin, is usually taken at home. Some are considered hazardous. There might be special precautions for storing and handling an immunotherapy drug. You might be told to be careful not to let others come into contact with it or your body fluids while taking it and for a time after taking it. Sometimes you need to wear gloves when touching the pills or capsules. Some drugs have to be kept in the bottle or box they came in. And some drugs and the packages they come in need to be disposed of in a certain way. Some might have to be taken back to the drug store to be thrown away safely. If you are taking an oral drug, talk to your cancer care team about whether special precautions are needed at home.

Keeping family and friends safe

Unless your health care team tells you differently, you can usually be around family and friends during the weeks and months you're getting immunotherapy. If you're getting treatment at a center, family and friends can often come with you. However, some treatment centers only allow patients in the infusion area and visitors may need to stay in the waiting room.

You are the only person who should be exposed to the drug you are getting, but any spilled IV drug, and any powder or dust from a pill or capsule, or any liquid from oral or topical immunotherapy might be hazardous to others if they are around it.

It's important to talk to your cancer care team and be aware of any special precautions that might be needed while you are taking an immunotherapy drug.

Hyperlinks

1. www.cancer.org/cancer/managing-cancer/treatment-types/chemotherapy.html

References

Brown VT. Targeted therapy. In Olsen MM, LeFebvre KB, Brassil KJ, eds. *Chemotherapy and Immunotherapy Guidelines and Recommendations for Practice*. Pittsburgh, PA: Oncology Nursing Society; 2019:103-139.

Last Revised: December 27, 2019

Written by

The American Cancer Society medical and editorial content team
(<https://www.cancer.org/cancer/acs-medical-content-and-news-staff.html>)

Our team is made up of doctors and oncology certified nurses with deep knowledge of cancer care as well as editors and translators with extensive experience in medical writing.

American Cancer Society medical information is copyrighted material. For reprint requests, please see our Content Usage Policy (www.cancer.org/about-us/policies/content-usage.html).

cancer.org | 1.800.227.2345