



[cancer.org](https://www.cancer.org) | 1.800.227.2345

Marijuana and Cancer

Marijuana is the name given to the dried buds and leaves of varieties of the *Cannabis sativa* plant, which can grow wild in warm and tropical climates throughout the world and be cultivated commercially. It goes by many names, including pot, grass, cannabis, weed, hemp, hash, marihuana, ganja, and dozens of others.

- [Types of marijuana compounds](#)
- [How can marijuana affect symptoms of cancer?](#)
- [Possible harmful effects of marijuana](#)
- [Cannabinoid drugs](#)
- [What does the American Cancer Society say about the use of marijuana in people with cancer?](#)

Marijuana has been used in herbal remedies for centuries. Scientists have identified many biologically active components in marijuana. These are called *cannabinoids*. The two best studied components are the chemicals *delta-9-tetrahydrocannabinol* (often referred to as THC), and *cannabidiol* (CBD). Other cannabinoids are being studied.

At this time, the US Drug Enforcement Administration (DEA) lists marijuana and its cannabinoids as Schedule I controlled substances. This means that they cannot legally be prescribed, possessed, or sold under federal law. Whole or crude marijuana (including marijuana oil or hemp oil) is not approved by the US Food and Drug Administration (FDA) for any medical use. But the use of marijuana to treat some medical conditions is legal under state laws in many states.

Dronabinol, a pharmaceutical form of THC, and a man-made cannabinoid drug called nabilone are approved by the FDA to treat some conditions.

Types of marijuana compounds

Different compounds in marijuana have different actions in the human body. For example, delta-9-tetrahydrocannabinol (THC) seems to cause the "high" reported by marijuana users, and also can help relieve [pain](#)¹ and [nausea](#)², reduce inflammation, and can act as an antioxidant. Cannabidiol (CBD) can help treat seizures, can reduce anxiety and paranoia, and can counteract the "high" caused by THC.

Different cultivars (strains or types) and even different crops of marijuana plants can have varying amounts of these and other active compounds. This means that marijuana can have different effects based on the strain used.

The effects of marijuana also vary depending on how marijuana compounds enter the body. The most common ways to use marijuana are in food (edible marijuana) and by smoking or vaping it (inhaled marijuana):

- **Edible marijuana:** When taken by mouth, such as when it's used in cooking oils, drinks (beer, tea, vodka, soda), baked goods (biscuits, brownies, cookies), and candy, the THC is absorbed poorly and can take hours to be absorbed. Once it's absorbed, it's processed by the liver, which produces a second psychoactive compound (a substance that acts on the brain and changes mood or consciousness) that affects the brain differently than THC. It's important to know that the amount of THC in foods that have had marijuana added to them is often unknown and getting too much THC might cause symptoms of overdose.
- **Inhaled marijuana:** When marijuana is smoked or vaporized, THC enters the bloodstream and goes to the brain quickly. The second psychoactive compound is produced in small amounts, and so has less effect. The effects of inhaled marijuana fade faster than marijuana taken by mouth.

How can marijuana affect symptoms of cancer?

A number of small studies of smoked marijuana found that it can be helpful in treating

Studies have long shown that people who took marijuana extracts in clinical trials tended to need less pain medicine.

More recently, scientists reported that THC and other cannabinoids such as CBD slow growth and/or cause death in certain types of cancer cells growing in lab dishes. Some animal studies also suggest certain cannabinoids may slow growth and reduce spread of some forms of cancer.

There have been some early clinical trials of cannabinoids in treating cancer in humans and more studies are planned. While the studies so far have shown that cannabinoids can be safe in treating cancer, they do not show that they help control or cure the disease.

Relying on marijuana alone as treatment while avoiding or delaying conventional medical care for cancer may have serious health consequences.

Possible harmful effects of marijuana

Marijuana can also pose some harms to users. While the most common effect of marijuana is a feeling of euphoria ("high"), it also can lower the user's control over movement, cause disorientation, and sometimes cause unpleasant thoughts or feelings of anxiety and paranoia.

Smoked marijuana delivers THC and other cannabinoids to the body, but it also delivers harmful substances to users and those close by, including many of the same substances found in tobacco smoke.

Because marijuana plants come in different strains with different levels of active compounds, it can make each user's experience very hard to predict. The effects can also differ

lower doses.

People who have had emotional illnesses, paranoia, or hallucinations may find their symptoms are worse when taking cannabinoid drugs.

Talk to your doctor about what you should expect when taking one of these drugs. It's a good idea to have someone with you when you first start taking one of these drugs and after any dose changes.

What does the American Cancer Society say about the use of marijuana in people with cancer?

The American Cancer Society supports the need for more scientific research on cannabinoids for cancer patients, and recognizes the need for better and more effective therapies that can overcome the often debilitating side effects of cancer and its treatment. The Society also believes that the classification of marijuana as a Schedule I controlled substance by the US Drug Enforcement Administration imposes nu/GSo8r.65 Tm 0 0 0 .qs6

4. www.cancer.org/cancer/managing-cancer/side-effects/pain/peripheral-neuropathy.html

References

Abrams DI, Jay CA, Shade SB, et al. Cannabis in painful HIV-associated sensory neuropathy: a randomized placebo-controlled trial. *Neurology*. 2007;68(7):515-521.

Ahmedzai S, Carlyle DL, Calder IT, Moran F. Anti-emetic efficacy and toxicity of nabilone, a synthetic cannabinoid, in lung cancer chemotherapy. *Br J Cancer*. 1983;48(5):657-663.

American College of Physicians. Supporting research into the therapeutic role of marijuana. (Position paper, 2008.) Accessed at www.acponline.org/advocacy/where_we_stand/other_issues/medmarijuana.pdf on March 4, 2015.

Bhattacharyya S, Crippa JA, Allen P, et al. Induction of psychosis by Δ^9 -tetrahydrocannabinol reflects modulation of prefrontal and striatal function during attentional salience processing. *Arch Gen Psychiatry*. 2012;69(1):27-36.

Beal JE, Olson R, Laubenstein L, et al. Dronabinol as a treatment for anorexia associated with weight loss in patients with AIDS. *J Pain Symptom Manage*. 1995;10(2):89-97.

Beal JE, Olson R, Lefkowitz L, et al. Long-term efficacy and safety of dronabinol for acquired immunodeficiency syndrome-associated anorexia. *J Pain Symptom Manage*. 1997;14(1):7-14.

Cannabis-In-Cachexia-Study-Group, Strasser F, Luftner D, Possinger K, et al. Comparison of orally administered cannabis extract and Δ^9 -tetrahydrocannabinol in treating patients with cancer-related anorexia-cachexia syndrome: a multicenter, phase III, randomized, double-blind, placebo-controlled clinical trial from the Cannabis-In-Cachexia-Study-Group. *J Clin Onc*. 2006;24:3394-3400.

Ellis RJ, Toperoff W, Vaida F, et al. Smoked medicinal cannabis for neuropathic pain in HIV: a randomized, crossover clinical trial. *Neuropsychopharmacology*. 2009;34(3):672-680.

Guzmán M, Duarte MJ, Blázquez C, et al. A pilot clinical study of Δ^9 -

Kramer JL. Medical marijuana for cancer. *CA Cancer J Clin*. 2014 Dec 10.

Meiri E, Jhangiani H, Vredenburg JJ, et al. Efficacy of dronabinol alone and in combination with ondansetron versus ondansetron alone for delayed chemotherapy-induced nausea and vomiting. *Curr Med Res Opin*. 2007;23(3):533-543.

Musty RE, Rossi R. Effects of Smoked Cannabis and Oral 9-Tetrahydrocannabinol on Nausea and Emesis After Cancer Chemotherapy: A Review of State Clinical Trials. *Journal of Cannabis Therapeutics*. 2001; 1(1): 29-56.

National Cancer Institute. Cannabis and Cannabinoids (PDQ®) Health Professional Version, updated 12/17/14. Accessed at www.cancer.gov/cancertopics/pdq/cam/cannabis/healthprofessional on March 4, 2015.

Portenoy RK, Ganae-Motan ED, Allende S, et al. Nabiximols for opioid-treated cancer patients with poorly-controlled chronic pain: a randomized, placebo-controlled, grad 0 gElj 0 JTs7t8th c

symptoms. *J Pain Symptom Manage*. 2005;29:358-367.

Last Revised: August 3, 2022

Written by

The American Cancer Society medical and editorial content team
([https](https://www.cancer.org)
