



[cancer.org](https://www.cancer.org) | 1.800.227.2345

Cervical Cancer Causes, Risk Factors, and Prevention

Learn about the risk factors for cervical cancer and what you might be able to do to help lower your risk.

Risk Factors

A risk factor is anything that affects your chance of getting a disease such as cancer. Learn more about the risk factors for cervical cancer.

- [Risk Factors for Cervical Cancer](#)
- [What Causes Cervical Cancer?](#)

Prevention

There is no way to completely prevent cancer. But there are things you can do that might lower your risk. Learn more.

- [Can Cervical Cancer Be Prevented?](#)

Risk Factors for Cervical Cancer

- [Risk factors you can possibly change](#)
- [Risk factors that cannot be changed](#)

- [Factors that may lower your risk](#)

A risk factor is anything that increases your chance of getting a disease such as cancer. Different cancers have different risk factors. For example, exposing skin to strong sunlight is a risk factor for skin cancer. Smoking is a risk factor for many cancers. But having a risk factor, or even several, does not mean that you will get the disease.

Several risk factors can increase your chance of developing cervical cancer. People without any of these risk factors rarely develop cervical cancer. Although these risk factors can increase the odds of developing cervical cancer, many with these risks do not develop this disease.

When you think about risk factors, it helps to focus on those you can change or avoid (like smoking or human papillomavirus infection), rather than those you cannot (such as your age and family history). However, it is still important to know about risk factors that cannot be changed, because it's even more important for those who have these factors to get regular screening tests to find cervical cancer early.

Risk factors you can possibly change

Human papillomavirus (HPV) infection

Infection by the [human papillomavirus](#)¹ (HPV) is the most important risk factor for cervical cancer. HPV is a group of more than 150 related viruses. Some of them cause a type of growth called **papillomas**, which are more commonly known as warts.

- HPV can infect cells on the surface of the skin, and those lining the genitals, anus, mouth and throat, but not the blood or internal organs such as the heart or lungs.
- HPV can spread from one person to another during skin-to-skin contact. One way HPV spreads is through sexual activity, including vaginal, anal, and even oral sex.
- Different types of HPV cause warts on different parts of the body. Some cause common warts on the hands and feet; others tend to cause warts on the lips or tongue.

Certain types of HPV may cause warts on or around the female and male genital organs and in the anal area. These are called low-risk types of HPV because they are seldom linked to cancer.

Other types of HPV are called **high-risk types** because they are strongly linked to

cancers, including cancer of the cervix, [vulva](#)², and [vagina](#)³ in women, [penile cancer](#)⁴ in men, and cancers of the [anus](#)⁵, [mouth, and throat](#)⁶ in both men and women.

Infection with HPV is common, and in most people the body can clear the infection by itself. Sometimes, however, the infection does not go away and becomes chronic. Chronic infection, especially when it is caused by certain high-risk HPV types, can eventually cause certain cancers, such as cervical cancer.

Although there is currently no cure for HPV infection, there are ways to treat the warts and abnormal cell growth that HPV causes. Also, [HPV vaccines](#)⁷ are available to help prevent infection by certain types of HPV and some of the cancers linked to those types.

For more information on this topic, see [HPV](#)⁸.

Sexual history

Several factors related to your sexual history can increase the risk of cervical cancer. The risk is most likely affected by increasing the chances of exposure to HPV.

- Becoming sexually active at a young age (especially younger than 18 years old)
- Having many sexual partners
- Having one partner who is considered high risk (someone with HPV infection or who has many sexual partners)

Smoking

When someone smokes, they and those around them are exposed to many cancer-causing chemicals that affect organs other than the lungs. These harmful substances are absorbed through the lungs and carried in the bloodstream throughout the body.

Women who smoke are about twice as likely as those who don't smoke to get cervical cancer. Tobacco by-products have been found in the cervical mucus of women who smoke. Researchers believe that these substances damage the DNA of cervix cells and may contribute to the development of cervical cancer. Smoking also makes the immune system less effective in fighting HPV infections.

Having a weakened immune system

[Human immunodeficiency virus \(HIV\)](#)⁹, the virus that causes AIDS, weakens the immune system and puts people at higher risk for HPV infections.

The immune system is important in destroying cancer cells and slowing their growth and spread. In women with HIV, a cervical pre-cancer might develop into an invasive cancer faster than it normally would.

Young age at first full-term pregnancy

Women who were younger than 20 years when they had their first full-term pregnancy are more likely to get cervical cancer later in life than women who waited to get pregnant until they were 25 years or older.

Economic status

Many low-income women do not have easy access to adequate health care services, including cervical cancer screening with Pap tests and HPV tests. This means they may not get screened or treated for cervical pre-cancers.

A diet low in fruits and vegetables

Women whose diets don't include enough fruits and vegetables may be at increased risk for cervical cancer.

Risk factors that cannot be changed

Diethylstilbestrol (DES) withGS155a4 0 g 0 0 .712 495.58 Tm /F2 12 Tf 0 0 0 g 0 01 72.6 75ee ca

9. www.cancer.org/cancer/risk-prevention/infections/hiv-infection-aids.html
10. www.cancer.org/cancer/risk-prevention/medical-treatments/des-exposure.html

References

- Adam E, Kaufman RH, Adler-Storthz K, et al. A prospective study of association of herpes simplex virus and human papillomavirus infection with cervical neoplasia in women exposed to diethylstilbestrol in utero. *Int J Cancer*. 1985;35:19-26.
- Castellsagué X, Díaz M, Vaccarella S, de Sanjosé S, Muñoz N, Herrero R, et al. Intrauterine device use, cervical infection with human papillomavirus, and risk of cervical cancer: A pooled analysis of 26 epidemiological studies. *Lancet Oncol*. 2011;12(11):1023.
- Centers for Disease Control and Prevention (CDC). Use of 9-Valent Human Papillomavirus (HPV) Vaccine: Updated HPV Vaccination Recommendations of the Advisory Committee on Immunization Practices. *MMWR Morb Mortal Wkly Rep*. 2015 March 27;64(11):300304.
- Chih HJ, Lee AH, Colville L, Binns CW, Xu D. A review of dietary prevention of human papillomavirus-related infection of the cervix and cervical intraepithelial neoplasia. *Nutr Cancer*. 2013;65:317–328.
- Cohen PA, Jhingran A, Oaknin A, Denny L. Cervical cancer. *Lancet*. 2019 Jan 12;393(10167):169-182. doi: 10.1016/S0140-6736(18)32470-X.
- Cortessis VK, Barrett M, Brown Wade N, Enebish T, Perrigo JL, Tobin J, et al. Intrauterine Device Use and Cervical Cancer Risk: A Systematic Review and Meta-analysis. *Obstet Gynecol*. 2017;130(6):1226.
- Eifel P, Klopp AH, Berek JS, and Konstantinopoulos A. Chapter 74: Cancer of the Cervix, Vagina, and Vulva. In: DeVita VT, Lawrence TS, Rosenberg SA, eds. DeVita, Hellman, and Rosenberg's *Cancer: Principles and Practice of Oncology*. 11th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2019.
- Fonseca-Moutinho JA. Smoking and cervical cancer. *ISRN Obstet Gynecol*. 2011;2011:847684. doi:10.5402/2011/847684.

Frumovitz M. Invasive cervical cancer: Epidemiology, risk factors, clinical manifestations, and diagnosis. UpToDate website. <https://www.uptodate.com/contents/invasive-cervical-cancer-epidemiology-risk-factors-clinical-manifestations-and-diagnosis>. Updated June 26, 2019. Accessed October 25, 2019.

Ghosh C, Baker JA, Moysich KB, et al. Dietary intakes of selected nutrients and food groups and risk of cervical cancer. *Nutr Cancer*. 2008;60:331-341.

Hatch EE, Herbst AL, Hoover RN, et al. Incidence of squamous neoplasia of the cervix and vagina in women exposed prenatally to diethylstilbestrol (United States). *Cancer Causes Control*. 2001;12:837-845.

Hernandez BY, Wilkens LR, Zhu X, et al. Transmission of human papillomavirus in heterosexual couples. *Emerg Infect Dis*. 2008;14(6):888-894.

Hogewoning CJ, Bleeker MC, van den Brule AJ, et al. Condom use promotes regression of cervical intraepithelial neoplasia and clearance of human papillomavirus: A randomized clinical trial. *Int J Cancer*. 2003;107(5):811-816.

Hoover RN, Hyer M, Pfeiffer RM, et al. Adverse health outcomes in women exposed in utero to diethylstilbestrol. *N Engl J Med*. 2011. 365: 1304-14.

International Collaboration of Epidemiological Studies of Cervical Cancer. Appleby P, Beral V, Berrington de González A, et al. Cervical cancer and hormonal contraceptives: Collaborative reanalysis of individual data for 16,573 women with cervical cancer and 35,509 women without cervical cancer from 24 epidemiological studies. *Lancet*. 2007;370:1609-1621.

International Collaboration of Epidemiological Studies of Cervical Cancer. Comparison of risk factors for invasive squamous cell carcinoma and adenocarcinoma of the cervix: collaborative reanalysis of individual data on 8,097 women with squamous cell carcinoma and 1,374 women with adenocarcinoma from 12 epidemiological studies. *Int J Cancer*. 2007 Feb 15;120(4):885-91.

Jhungrán A, Russell AH, Seiden MV, Duska LR, Goodman A, Lee S, et al. Chapter 84: Cancers of the Cervix, Vulva, and Vagina. In: Niederhuber JE, Armitage JO, Doroshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa: Elsevier; 2020.

Lu B, Wu Y, Nielson CM, et al. Factors associated with acquisition and clearance of

human papillomavirus infection in a cohort of US men: a prospective study. *J Infect Dis.* 2009;199(3):362371.

National Cancer Institute. Physician Data Query (PDQ). Cervical Cancer Treatment – Health Professional Version. 2019. <https://www.cancer.gov/types/cervical/hp/cervical-treatment-pdq>. Updated February 6, 2019. Accessed on September 5, 2019.

National Comprehensive Cancer Network (NCCN). Clinical Practice Guidelines in Oncology: Cervical Cancer. Version 4.2019. Accessed at https://www.nccn.org/professionals/physician_gls/pdf/cervical.pdf on September 5, 2019.

Roura E, Castellsagué X, Pawlita M, et al. Smoking as a major risk factor for cervical cancer and pre-cancer: Results from the EPIC cohort. *Int J Cancer.* 2014; 135: 453–66.

Schiffman M, Castle PE, Jeronimo J, et al. Human papillomavirus and cervical cancer. *Lancet.* 2007;370(9590):890907.

Silva J, Cerqueira F, Medeiros R. Chlamydia trachomatis infection: implications for HPV status and cervical cancer. *Arch Gynecol Obstet.* 2014 Apr;289(4):715-23. doi: 10.1007/s00404-013-3122-3.

Tokudome S, Suzuki S, Ichikawa H, et al. Cancer risk in women prenatally exposed to diethylstilbestrol. *Int J Cancer.* 2007;121:356-360.

Winer RL, Hughes JP, Feng Q, et al. Condom use and the risk of genital human papillomavirus infection in young women. *N Engl J Med.* 2006;354:26452654.

Winer RL, Lee SK, Hughes JP, et al. Genital human papillomavirus infection: incidence and risk factors in a cohort of female university students. *Am J Epidemiol.* 2003;157(3):218-226. Erratum in: *Am J Epidemiol.* 2003;157(9):858.

Zhu H, Shen Z, Luo H, Zhang W, Zhu X. Chlamydia Trachomatis Infection-Associated Risk of Cervical Cancer: A Meta-Analysis. *Medicine (Baltimore).* 2016;95(13):e3077. doi:10.1097/MD.0000000000003077

Last Revised: January 3, 2020

What Causes Cervical Cancer?

In recent years, there has been a lot of progress in understanding what happens in cells of the cervix when cancer develops. In addition, several risk factors have been identified that increase the odds that a woman might develop cervical cancer (see [Risk Factors for Cervical Cancer](#)).

The development of normal human cells mostly depends on the information contained in the cells' DNA. DNA is the chemical in our cells that makes up our genes, which control how our cells work. We look like our parents because they are the source of our DNA. But DNA affects more than just how we look.

Some genes control when cells grow, divide, and die:-

- Genes that help cells grow, divide, and stay alive are called **oncogenes**.
- Genes that help keep cell growth under control or make cells die at the right time are called **tumor suppressor genes**.

Cancers can be caused by DNA mutations (gene defects) that turn on oncogenes or turn off tumor suppressor genes.

[Human papillomaviruses](#)¹ (HPV) have two proteins known as E6 and E7 which turn off some tumor suppressor genes, such as p53 and Rb. This may allow the cells lining the cervix to grow too much and to develop changes in additional genes, which in some cases can lead to cancer.

But HPV is not the only cause of cervical cancer. Most women with HPV don't get cervical cancer, and other risk factors, like smoking and HIV infection, influence which women exposed to HPV are more likely to develop cervical cancer.

Hyperlinks

1. www.cancer.org/cancer/risk-prevention/hpv.html

References

Jhungran A, Russell AH, Seiden MV, Duska LR, Goodman A, Lee S, et al. Chapter 84: Cancers of the Cervix, Vulva, and Vagina. In: Niederhuber JE, Armitage JO, Doroshow

JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa: Elsevier; 2020.

Yim EK, Park JS. The role of HPV E6 and E7 oncoproteins in HPV-associated cervical carcinogenesis. *Cancer Res Treat*. 2005;37(6):319–324. doi:10.4143/crt.2005.37.6.319.

Last Revised: January 3, 2020

Can Cervical Cancer Be Prevented?

- [Finding cervical pre-cancers](#)
- [Things to do to prevent pre-cancers and cancers](#)

The two most important things you can do to prevent cervical cancer are to get the HPV vaccine if you are eligible, and to be tested regularly according to American Cancer Society (ACS) guidelines. These can be found in [The American Cancer Society Guidelines for the Prevention and Early Detection of Cervical Cancer](#)¹.

The most common form of cervical cancer starts with pre-cancerous changes and there are ways to stop this from developing. The first way is to find and treat pre-cancers before they become invasive cancers, and the second is to prevent the pre-cancers.

Finding cervical pre-cancers

A well-proven way to prevent cervical cancer is to have screening tests. Screening is having tests to find conditions that may lead to cancers and can find pre-cancers before they can turn into invasive cancer. The Pap test (or Pap smear) and the human0 gintacpv4(Things to c

- HPV vaccination of children between the ages of 9 and 12.
- Children and young adults age 13 through 26 who have not been vaccinated, or who haven't gotten all their doses, should get the vaccine as soon as possible. Vaccination of young adults will not prevent as many cancers as vaccination of children and teens.
- The ACS does not recommend HPV vaccination for persons older than 26 years.

It's important to know that no vaccine provides complete protection against all cancer-causing types of HPV, so routine cervical cancer screening is still needed.

For more information on the vaccine and HPV, please see [HPV Vaccines⁴](#).

Limit exposure to HPV

HPV is passed from one person to another during skin-to-skin contact with an infected area of the body. Although HPV can be spread during skin to skin contact including vaginal, anal, and oral sex sex doesn't have to occur for the infection to spread. All that is needed is skin-to-skin contact with an area of the body infected with HPV. This means that the virus can be spread without sex. It is even possible for a genital infection to spread through hand-to-genital contact.

Also, HPV infection seems to be able to spread from one part of the body to another. This means that an infection may start in the cervix and then spread to the vagina and vulva.

It can be very hard not to be exposed to HPV. It may be possible to prevent HPV infection by not allowing others to have contact with your anal or genital area, but even then there might be other ways to become infected that aren't yet clear.

Limiting the number of sex partners and avoiding sex with people who have had many other sex partners may lower your risk of exposure to HPV. But again, HPV is very common, so having sexual activity with even one other person can put you at risk. Remember that someone can have HPV for years and still have no symptoms. So it's possible someone can have the virus and pass it on without knowing it.

Use a condom

Condoms ("rubbers") provide some protection against HPV but they don't completely prevent infection. One reason that condoms cannot protect completely is because they don't cover every possible HPV-infected area of the body, such as skin of the genital or

Group. Smoking and cervical cancer: Pooled analysis of the IARC multi-centric case-control study. *Cancer Causes Control*. 2003;14:805-814.

Roura E, Castellsagué X, Pawlita M, et al. Smoking as a major risk factor for cervical cancer and pre-cancer: Results from the EPIC cohort. *Int J Cancer*. 2014; 135: 453–66.

Saslow D, Andrews KS, Manassaram-Baptiste D, et al. Human papillomavirus vaccination 2020 guideline update: American Cancer Society guideline adaptation. *CA Cancer J Clin*