



cancer.org | 1.800.227.2345

Lung Cancer Causes, Risk Factors, and Prevention

Learn about the risk factors for lung cancer and what you might be able to do to help lower your risk.

Risk Factors

A risk factor is anything that affects your chance of getting a disease such as cancer. Learn more about the risk factors for lung cancer.

- [Lung Cancer Risk Factors](#)
- [What Causes Lung Cancer?](#)

Prevention

There is no way to completely prevent cancer. But there are things you can do that might lower your risk. Learn more.

- [Can Lung Cancer Be Prevented?](#)

Lung Cancer Risk Factors

A risk factor is anything that increases a person's chance of getting a disease such as cancer.

If you or someone you care about needs help quitting, see [How to Quit Using Tobacco](#)⁴ or call the American Cancer Society at 1-800-227-2345

Exposure to radon

Radon is a naturally occurring radioactive gas that results from the breakdown of uranium in soil and rocks. You can't see, taste, or smell it. According to the US Environmental Protection Agency (EPA), radon is the second-leading cause of lung cancer in the United States, and it's the leading cause among people who don't smoke.

Outdoors, there is so little radon that it is not likely to be dangerous. But indoors, radon can be more concentrated. Breathing it in exposes your lungs to small amounts of radon. This may increase a person's risk of lung cancer.

Homes and other buildings in nearly any part of the country can have high indoor radon levels (especially in basements).

For more information, see [Radon and Cancer](#)⁵.

Exposure to asbestos

People who work with asbestos (such as in mines, mills, textile plants, places where insulation is used, and shipyards) are several times more likely to die of lung cancer. Lung cancer risk is much greater in workers exposed to asbestos who also smoke. It's not clear how much low-level or short-term exposure to asbestos might raise lung cancer risk.

- Radioactive ores, such as uranium
- Inhaled chemicals, such as arsenic, beryllium, cadmium, silica, vinyl chloride, nickel compounds, chromium compounds, coal products, mustard gas, and chloromethyl ethers
- [Diesel exhaust](#)⁸

The government and industry have taken steps in recent years to help protect workers from many of these exposures. But the dangers are still there, so if you work around these agents, be careful to limit your exposure whenever possible.

Taking certain dietary supplements

Studies looking at the possible role of vitamin supplements in reducing lung cancer risk have had disappointing results. In fact, multiple studies found that people who smoked who took beta-carotene supplements actually had an increased risk of lung cancer. The results of these studies suggest that people should avoid taking beta-carotene supplements.

Arsenic in drinking water

Studies of people in parts of Southeast Asia and South America with high levels of [arsenic](#)⁹ in their drinking water have found a higher risk of lung cancer. In most of these studies, the levels of arsenic in the water were many times higher than those typically seen in the United States, even areas where arsenic levels are above normal. For most Americans who are on public water systems, drinking water is not a major source of arsenic.

Risk factors you cannot change

Previous radiation therapy to the lungs



cancer slightly. This risk is far less than the risk caused by smoking, but about 1% to 2% of all deaths from lung cancer in the United States are thought to be due to outdoor air pollution.

Personal or family history of lung cancer

If you have had lung cancer, you have a higher risk of developing another lung cancer.

Brothers, sisters, and children of people who have had lung cancer may have a slightly higher risk of lung cancer themselves, especially if the relative was diagnosed at a younger age. It's not clear how much of this risk might be due to shared genes among family members and how much might be from shared household exposures (such as

The Alpha-Tocopherol, Beta Carotene Cancer Prevention Study Group. The effect of vitamin E and beta carotene on the incidence of lung cancer and other cancers in male smokers. *N Engl J Med*. 1994;330:1029-1035.

Amos CI, Pinney SM, Li Y, et al. A susceptibility locus on chromosome 6q greatly increases lung cancer risk among light and never smokers. *Cancer Res*. 2010;70:2359–2367.

Bein K, Leikauf GD. Acrolein - a pulmonary hazard. *Mol Nutr Food Res*. 2011 Sep;55(9):1342-60. doi: 10.1002/mnfr.201100279. PMID: 21994168.

Blot WJ, Cohen SS, Aldrich M, McLaughlin JK, Hargreaves MK, Signorello LB. Lung cancer risk among smokers of menthol cigarettes. *J Natl Cancer Inst*. 2011 May 18;103(10):810-6. doi: 10.1093/jnci/djr102. Epub 2011 Mar 23. PMID: 21436064; PMCID: PMC3096798.

Callaghan RC, Allebeck P and Sidorchuk A. Marijuana use and risk of lung cancer: a 40-year cohort study. *Cancer Causes Control*. 2013 Oct;24(10):1811-20.

Environmental Protection Agency. Health Risk of Radon.
<https://www.epa.gov/radon/health-risk-radon>. Last updated 01/09/2024. Accessed Jan 23, 2024.

Omenn GS, Goodman GE, Thornquist et al. Risk factors for lung cancer and for intervention effects in CARET, the Beta-Carotene and Retinol Efficacy Trial. *J Natl Cancer Inst*. 1996;88:1550-1559.

Owen KP, Sutter ME, Albertson TE. Marijuana: respiratory tract effects. *Clin Rev Allergy Immunol*. 2014 Feb;46(1):65-81. doi: 10.1007/s12016-013-8374-y. PMID: 23715638.

Pallis AG, Syrigos KN. Lung cancer in never smokers: disease characteristics and risk factors. *Crit Rev Oncol Hematol*. 2013 Dec;88(3):494-503. doi: 10.1016/j.critrevonc.2013.06.011. Epub 2013 Aug 4. PMID: 23921082.

Schabath MB, Cote ML. Cancer Progress and Priorities: Lung Cancer. *Cancer Epidemiol Biomarkers Prev*. 2019 Oct;28(10):1563-1579. doi: 10.1158/1055-9965.EPI-19-0221. PMID: 31575553; PMCID: PMC6777859.

Tashkin DP. Effects of marijuana smoking on the lung. *Ann Am Thorac Soc*. 2013 Jun;10(3):239-47. doi: 10.1513/AnnalsATS.201212-127FR. PMID: 23802821.

US Department of Health and Human Services. *The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General, 2014*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2014.

Last Revised: January 29, 2024

What Causes Lung Cancer?

We don't know what causes each case of lung cancer. But we do know many of the risk factors for these cancers (see [Lung Cancer Risk Factors](#)) and how some of them cause normal cells to become cancer cells.

- [How smoking leads to lung cancer](#)
- [Causes in people who don't smoke](#)
- [Gene changes that may lead to lung cancer](#)
- [Inherited gene changes \(germline mutations\)](#)

How smoking leads to lung cancer

[Smoking tobacco](#)¹ is by far the leading cause of lung cancer. About 80% of lung cancer deaths are caused by smoking, and many others are caused by exposure to secondhand smoke.

Smoking is clearly the strongest risk factor for lung cancer, but it often interacts with other factors. People who smoke and are exposed to other known risk factors, such as [radon](#)² and [asbestos](#)³, are at an even higher risk. Not everyone who smokes gets lung cancer, so other factors like genetics probably play a role as well (see below).

Causes in people who don't smoke

Not all people who get lung cancer smoke. Many people with lung cancer formerly smoked, but many others never smoked at all.

Lung cancer in people who don't smoke can be caused by exposure to radon,

were at an increased hereditary risk for developing lung cancer, but it could also help guide discussion about the best way to treat your lung cancer. If you have lung cancer and are also found to have a certain germline mutation, you may respond well to that mutation's targeted therapy.

Examples of possible germline mutations for patients with lung cancer include: CHEK2, ATM, TP53, *BRCA1*, EGFR, APC, and PALB2. Studies are ongoing to better understand the role of germline mutations in lung cancer. Regardless of whether you carry a higher hereditary risk for lung cancer, doctors recommend that all people avoid tobacco smoke and other exposures that will increase cancer risk.

Acquired gene changes (somatic mutations or “driver mutations”)

Acquired gene changes, or somatic mutations, may occur in any individual cell and cannot be inherited. Somatic mutations refer to DNA changes within cells that were not passed from your parents, but rather were acquired during your lifetime. Certain somatic mutations can affect the cell's ability to control its own growth, and will eventually transform a noncancer cell to become a cancer cell. These somatic mutations are also known as “driver mutations.” If your tumor is found to have a driver mutation, you will likely respond well to targeted therapy.

For patients with advanced non-small cell lung cancer, it is recommended that the lung mass or a metastatic mass be tested for driver mutations. It is standard practice to test for the following driver mutations: EGFR, ALK, ROS1, MET, RET, BRAF, and NTRK. If any of these driver mutations are found, initial treatment with a targeted therapy (rather than chemotherapy) would be recommended. Although these mutations can potentially be found in any patient with lung cancer, nonsmokers with lung cancer are more likely to have a driver mutation.

Hyperlinks

1. www.cancer.org/cancer/risk-prevention/tobacco.html
2. www.cancer.org/cancer/risk-prevention/radiation-exposure/radon.html
3. www.cancer.org/cancer/risk-prevention/chemicals/asbestos.html
4. www.cancer.org/cancer/risk-prevention/tobacco/health-risks-of-tobacco/secondhand-smoke.html
5. www.cancer.org/cancer/risk-prevention/chemicals/diesel-exhaust-and-cancer.html

References

Amos CI, Pinney SM, Li Y, et al. A susceptibility locus on chromosome 6q greatly increases lung cancer risk among light and never smokers. *Cancer Res*. 2010;70:2359–2367.

Araujo LH, Horn L, Merritt RE, Shilo K, Xu-Welliver M, Carbone DP. Ch. 69 - Cancer of the Lung: Non-small cell lung cancer and small cell lung cancer. In: Niederhuber JE, Armitage JO, Doroshow JH, Kastan MB, Tepper JE, eds. *Abeloff's Clinical Oncology*. 6th ed. Philadelphia, Pa: Elsevier; 2020.

Chiang A, Detterbeck FC, Stewart T, Decker RH, Tanoue L. Chapter 48: Non-small cell lung cancer. In: DeVita VT, Lawrence TS, Rosenberg SA, eds. *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology*. 11th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2019.

Gazdar AF. Activating and resistance mutations of EGFR in non-small-cell lung cancer: role in clinical response to EGFR tyrosine kinase inhibitors. *Oncogene*. 2009;28 Suppl 1(Suppl 1):S24–S31.

Hann CL, Wu A, Rekhtman N, Rudin CM. Chapter 49: Small cell and Neuroendocrine Tumors of the Lung. In: DeVita VT, Lawrence TS, Rosenberg SA, eds. *DeVita, Hellman, and Rosenberg's Cancer: Principles and Practice of Oncology*. 11th ed. Philadelphia, Pa: Lippincott Williams & Wilkins; 2019.

National Cancer Institute. Physician Data Query (PDQ). Patient Version. Non-Small Cell Lung Cancer Treatment. 2023. Accessed at <https://www.cancer.gov/types/lung/patient/non-small-cell-lung-treatment-pdq> on Jan 24, 2024.

Sorscher S, LoPiccolo J, Chen E, et al. Landscape of pathogenic germline variants in patients with lung cancer. Presented at ASCO Plenary Series; August 16, 2022. Abstract 388570.

Varella-Garcia M. Chromosomal and genomic changes in lung cancer. *Cell Adh Migr*. 2010;4(1):100–106.

Last Revised: January 29, 2024

Can Lung Cancer Be Prevented?

Not all lung cancers can be prevented, but you may be able to lower your risk for lung cancer by changing the [risk factors](#) that you can control.

- [Stay away from tobacco](#)
- [Avoid radon exposure](#)
- [Avoid or limit exposure to cancer-causing agents](#)
- [Eat a healthy diet](#)

Stay away from tobacco

The best way to reduce your risk of lung cancer is not to smoke and to avoid breathing in other people's smoke.

If you stop smoking before a cancer develops, your damaged lung tissue gradually starts to repair itself. No matter what your age or how long you've smoked, quitting will lower your risk of lung cancer and help you live longer. If you would like help quitting smoking, see [How to Quit Using Tobacco](#)¹ or call the American Cancer Society at 1-800-227-2345.

Avoid radon exposure

Trying to reduce the risk of lung cancer in people who currently smoke or those who formerly smoked by giving them high doses of vitamins or vitamin-like drugs has not

Our team is made up of doctors and oncology certified nurses with deep knowledge of cancer care as well as journalists, editors, and translators with extensive experience in medical writing.

American Cancer Society medical information is copyrighted material. For reprint requests, please see our Content Usage Policy (www.cancer.org/about-us/policies/content-usage.html).

cancer.org | 1.800.227.2345