



[cancer.org](http://cancer.org) | 1.800.227.2345

---

# Vulvar Cancer Causes, Risk Factors, and Prevention

Learn about the risk factors for vulvar cancer and what you might be able to do to help lower your risk.

## Risk Factors

A risk factor is anything that affects your chance of getting a disease such as cancer. Learn more about the risk factors for vulvar cancer.

- [Risk Factors for Vulvar Cancer](#)
- [What Causes Vulvar Cancer?](#)

## Prevention

There is no way to completely prevent cancer. But there are things you can do that might lower your risk. Learn more.

- [Can Vulvar Cancer Be Prevented?](#)

---

# Risk Factors for Vulvar Cancer

- [Age](#)
- [Human papillomavirus](#)

- [Smoking](#)
- [HIV infection](#)
- [Vulvar intraepithelial neoplasia \(VIN\)](#)
- [Lichen sclerosus](#)
- [Other genital cancers](#)
- [Melanoma or atypical moles](#)

A risk factor is anything that changes a person's chance of getting a disease such as cancer. Different cancers have different risk factors. For example, exposing skin to strong sunlight is a risk factor for skin cancer. Smoking is a risk factor for many cancers.

There are different kinds of risk factors. Some, such as your age or race, can't be changed. Others may be related to personal choices such as smoking, drinking, or diet. Some factors influence risk more than others. But risk factors don't tell us everything. Having a risk factor, or even several, does not mean that a person will get the disease. Also, not having any risk factors doesn't mean that you won't get it, either.

Although several risk factors increase the odds of developing vulvar cancer, most women with these risks do not develop it. And some women who don't have any apparent risk factors develop vulvar cancer. When a woman develops vulvar cancer, it is usually not possible to say with certainty that a particular risk factor was the cause.

## **Age**

The risk of vulvar cancer goes up as women age. Less than 20% of cases are in women younger than age 50, and more than half occur in women over age 70. The average

several inches across. The medical term for genital warts is **condyloma acuminatum**. Two types of HPV (HPV 6 and HPV 11) cause most cases of genital warts, but are seldom linked to cancer and are known as **low-risk** HPV.

Other HPV types have been linked with cancers of the [cervix](#)<sup>1</sup>, [vagina](#)<sup>2</sup>, and vulva in women, cancer of the [penis](#)<sup>3</sup> in men, and cancers of the anus and [throat](#)<sup>4</sup> (in men and women). These are known as **high-risk** types of HPV and include HPV 16 and HPV 18 as well as others. Infection with a high-risk HPV may produce no visible signs until pre-cancerous changes or cancer develops.

HPV can pass from one person to another during skin-to-skin contact. One way HPV is spread is through sexual activity, including vaginal and anal intercourse and even oral sex.

Some doctors think there are 2 kinds of vulvar cancer. One kind is associated with HPV infection (more than half of all vulvar cancers are linked to infection with the high-risk HPV types) and tends to occur in younger women. The other is not associated with HPV infection, is more often found in older women, and may develop from a precursor lesion called **differentiated vulvar intraepithelial neoplasia** (discussed below).

Vaccines have been developed to help prevent infection with some types of HPV.

For more about HPV, see [HPV \(Human Papillomavirus\)](#)<sup>5</sup>.

## Smoking

Smoking exposes people to many cancer-causing chemicals that affect more than their lungs. These harmful substances can be absorbed into the lining of the lungs and spread throughout the body. Smoking increases the risk of developing vulvar cancer. Among women who have a history of HPV infection, smoking further increases the risk of developing vulvar cancer. If women are infected with a high-risk HPV, they have a much higher risk of developing vulvar cancer if they smoke.

## HIV infection

HIV (human immunodeficiency virus) causes AIDS (acquired immunodeficiency syndrome). Because this virus damages the immune system, it makes women more likely to get and to stay infected with HPV. This could increase the risk of vulvar pre-cancer and cancer. Scientists also believe that the immune system plays a role in destroying cancer cells and slowing their growth and spread.

## **Vulvar intraepithelial neoplasia (VIN)**

Squamous cell carcinoma of the vulva usually forms slowly over many years. Pre-cancerous changes often occur first and can last for several years. The medical term most often used for this pre-cancerous condition is **vulvar intraepithelial neoplasia**

by LSA, with about 4% of women having LSA later developing vulvar cancer.

## Other genital cancers

Women with [cervical cancer](#)<sup>6</sup> also have a higher risk of vulvar cancer. This is probably because these cancers share certain risk factors. The same HPV types that are linked to cervical cancer are also linked to vulvar cancer. Smoking is also linked to a higher risk of both cervical and vulvar cancers.

## Melanoma or atypical moles

Women who have had [melanoma](#)<sup>7</sup> or dysplastic nevi (atypical moles) in other places have an increased risk of developing a melanoma on the vulva. A family history of melanoma also leads to an increased risk.

## Hyperlinks

1. [www.cancer.org/cancer/types/cervical-cancer.html](http://www.cancer.org/cancer/types/cervical-cancer.html)
2. [www.cancer.org/cancer/types/vaginal-cancer.html](http://www.cancer.org/cancer/types/vaginal-cancer.html)
3. [www.cancer.org/cancer/types/penile-cancer.html](http://www.cancer.org/cancer/types/penile-cancer.html)
4. [www.cancer.org/cancer/types/oral-cavity-and-oropharyngeal-cancer.html](http://www.cancer.org/cancer/types/oral-cavity-and-oropharyngeal-cancer.html)
5. [www.cancer.org/cancer/risk-prevention/hpv.html](http://www.cancer.org/cancer/risk-prevention/hpv.html)
6. [www.cancer.org/cancer/types/cervical-cancer.html](http://www.cancer.org/cancer/types/cervical-cancer.html)
7. [www.cancer.org/cancer/types/melanoma-skin-cancer.html](http://www.cancer.org/cancer/types/melanoma-skin-cancer.html)
8. [www.cancer.org/cancer/types/vulvar-cancer/references.html](http://www.cancer.org/cancer/types/vulvar-cancer/references.html)

## References

[See all references for Vulvar Cancer](#)

Last Revised: January 16, 2018

# What Causes Vulvar Cancer?

Several [risk factors](#) for cancer of the vulva have been identified, and we are beginning to understand how these factors can cause cells in the vulva to become cancerous.

Researchers have made a lot of progress in understanding how certain changes in DNA can cause normal cells to become cancerous. DNA is the chemical that carries the instructions for nearly everything our cells do. We usually look like our parents because they are the source of our DNA. However, DNA affects more than our outward appearance. Some genes (parts of our DNA) contain instructions for controlling when our cells grow and divide.

- Certain genes that promote cell division are called **oncogenes**.
- Others that slow down cell division or cause cells to die at the right time are called **tumor suppressor genes**.

Cancers can be caused by DNA mutations (defects) that turn on oncogenes or turn off tumor suppressor genes. Usually DNA mutations related to cancers of the vulva occur during life rather than having been inherited before birth. Acquired mutations may result from cancer-causing chemicals in tobacco smoke. Sometimes they occur for no apparent reason. For more on genes and cancer, see [Oncogenes and Tumor Suppressor Genes](#)<sup>1</sup>.

Studies suggest that squamous cell cancer of the vulva (the most common type) can develop in at least 2 ways. In up to half of cases, human papillomavirus (HPV) infection appears to have an important role. Vulvar cancers associated with HPV infection (the basaloid and warty subtypes) seem to have certain distinctive features. They are often found along with several other areas of vulvar intraepithelial neoplasia (VIN). The women who have these cancers tend to be younger and often smoke.

The second process by which vulvar cancers develop does not involve HPV infection. Vulvar cancers not linked to HPV infection (the keratinizing subtype) are usually diagnosed in older women (over age 55). These women may have lichen sclerosis and may also have the differentiated type of VIN. DNA tests from vulvar cancers in older women rarely show HPV infection, but often show mutations of the *p53* tumor suppressor gene. The *p53*

prevent cancer of the vulva and at some point might lead to changes in treatment.

Because vulvar melanomas and adenocarcinomas are so rare, much less is known about how they develop.

## Hyperlinks

1. [www.cancer.org/cancer/understanding-cancer/genes-and-cancer/oncogenes-tumor-suppressor-genes.html](http://www.cancer.org/cancer/understanding-cancer/genes-and-cancer/oncogenes-tumor-suppressor-genes.html)
2. [www.cancer.org/cancer/types/vulvar-cancer/references.html](http://www.cancer.org/cancer/types/vulvar-cancer/references.html)

## References

[See all references for Vulvar Cancer](#)

Last Revised: January 16, 2018

---

# Can Vulvar Cancer Be Prevented?

- [Avoid HPV infection](#)
- [Get vaccinated](#)
- [Don't smoke](#)
- [Get regular pelvic checkups](#)

The risk of vulvar cancer can be lowered by avoiding certain [risk factors](#) and by having pre-cancerous conditions [treated](#) before an invasive cancer develops. Taking these steps cannot guarantee that all vulvar cancers are prevented, but they can greatly reduce your chances of developing vulvar cancer.

## Avoid HPV infection

Infection with human papillomavirus (HPV) is a risk factor for vulvar cancer. In women,

HPV infections occur mainly at younger ages and are less common in women over 30. The reason for this is not clear.

HPV is passed from one person to another during skin-to-skin contact with an infected area of the body. HPV can be spread during sexual activity -- including vaginal intercourse, anal intercourse, and oral sex -- but sex doesn't have to occur for the infection to spread. All that is needed is skin-to-skin contact with an area of the body infected with HPV. The virus can be spread through genital-to-genital contact. It is even possible to spread a genital infection through hand-to-genital contact.

An HPV infection also seems to be able to be spread from one part of the body to another. This means that an infection may start in the cervix and then spread to the vagina and vulva.

It can be very hard to avoid being exposed to HPV. If you are sexually active, limiting the number of sex partners and avoiding sex with people who have had many other sex partners can help lower your risk of exposure to HPV. But again, HPV is very common, so having sex with even one other person can put you at risk.

Infection with HPV is common, and in most cases your body is able to clear the infection on its own. But in some cases, the infection does not go away and becomes chronic. Chronic infection, especially with high-risk HPV types, can eventually cause certain cancers, including vulvar cancer.

A person can be infected with HPV for years and not have any symptoms, so the absence of visible warts cannot be used to tell if someone has HPV. Even when someone doesn't have warts (or any other symptom), he (or she) can still be infected with HPV and pass the virus to somebody else.

## **Condoms**

Condoms (rubbers) provide some protection against HPV, but they do not completely prevent infection. Condoms cannot protect completely because they don't cover every possible HPV-infected area of the body, such as the skin on the genital or anal area. Still, condoms do provide some protection against HPV, and they also protect against HIV and some other sexually transmitted diseases.

## **Get vaccinated**

Vaccines that protect against certain HPV infections are available. All of them protect against infection with HPV subtypes 16 and 18. Some can also protect against





for any visible abnormalities. The health care professional then places a speculum inside the vagina. A speculum is a metal or plastic instrument that keeps the vagina open so that the cervix can be seen clearly. If a Pap test or HPV test are being done during a pelvic exam, a sample of cells and mucus is lightly scraped from the exocervix (the surface of the cervix that is closest to the vagina) and the endocervix (the inside part of the cervix that is closest to the body of the uterus). Then, the speculum is removed. The doctor then will check the organs of the pelvis by inserting 1 or 2 gloved fingers of one hand into the vagina while the lower abdomen is palpated (felt), just above the pubic bone, with the other hand. The doctor may do a rectal exam at this time also. It is very important to know that a Pap test and HPV test are not always done when a pelvic exam is done, so if you are uncertain, you should ask if one was done.

### **Self-exam of the vulva**

For most women, the best way to find VIN and vulvar cancer is to report any signs and symptoms to their health care provider and have a yearly well-woman exam. If you have an increased risk of vulvar cancer, you may also want to check your vulva regularly to look for any of the signs of vulvar cancer. This is known as self-examination. Some women choose to examine themselves monthly using a mirror. This can allow you to become aware of any changes in the skin of your vulva. If you do this, look for any areas that are white, darkly pigmented, or red and irritated. You should also note any new growths, nodules, bumps, or ulcers (open sores). Report any of these to a doctor, since they could indicate a vulvar cancer or pre-cancer.

### **Hyperlinks**

1. [www.cancer.org/cancer/risk-prevention/hpv/hpv-vaccines.html](http://www.cancer.org/cancer/risk-prevention/hpv/hpv-vaccines.html)
2. [www.cancer.org/cancer/risk-prevention/hpv.html](http://www.cancer.org/cancer/risk-prevention/hpv.html)
3. [www.cancer.org/cancer/types/cervical-cancer.html](http://www.cancer.org/cancer/types/cervical-cancer.html)
4. [www.cancer.org/cancer/types/vulvar-cancer/references.html](http://www.cancer.org/cancer/types/vulvar-cancer/references.html)

### **References**

[See all references for Vulvar Cancer](#)

Last Revised: July 21, 2020

**Written by**

The American Cancer Society medical and editorial content team  
(<https://www.cancer.org/cancer/acs-medical-content-and-news-staff.html>)

Our team is made up of doctors and oncology certified nurses with deep knowledge of cancer care as well as journalists, editors, and translators with extensive experience in medical writing.

American Cancer Society medical information is copyrighted material. For reprint requests, please see our Content Usage Policy ([www.cancer.org/about-us/policies/content-usage.html](http://www.cancer.org/about-us/policies/content-usage.html)).

**cancer.org | 1.800.227.2345**